



The Common Follicle-Stimulating Hormone Receptor (FSHR) Promoter Polymorphism FSHR -29G>A Affects Androgen Production in Normal Human Small Antral Follicles

Borgbo, Tanni Kjær; Kluková, Hana; Macek, Milan; Chrudimska, Jana; Kristensen, Stine Gry; Hansen, Lise Lotte; Andersen, Claus Yding

Published in:
Frontiers in Endocrinology

DOI:
[10.3389/fendo.2017.00122](https://doi.org/10.3389/fendo.2017.00122)

Publication date:
2017

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Borgbo, T. K., Kluková, H., Macek, M., Chrudimska, J., Kristensen, S. G., Hansen, L. L., & Andersen, C. Y. (2017). The Common Follicle-Stimulating Hormone Receptor (FSHR) Promoter Polymorphism FSHR -29G>A Affects Androgen Production in Normal Human Small Antral Follicles. *Frontiers in Endocrinology*, 8, [122]. <https://doi.org/10.3389/fendo.2017.00122>



The Common Follicle-Stimulating Hormone Receptor (FSHR) Promoter Polymorphism FSHR –29G > A Affects Androgen Production in Normal Human Small Antral Follicles

Tanni Borgbo^{1,2}, Hana Klučková³, Milan Macek Sr.³, Jana Chrudimska³, Stine Gry Kristensen¹, Lise Lotte Hansen² and Claus Yding Andersen^{1*}

¹Laboratory of Reproductive Biology, The Juliane Marie Centre for Women, Children and Reproduction, University Hospital of Copenhagen, Copenhagen, Denmark, ²Department of Biomedicine, University of Aarhus, Aarhus, Denmark, ³Department of Biology and Medical Genetics, 2nd Faculty of Medicine Charles University, University Hospital Motol, Prague, Czechia

OPEN ACCESS

Edited by:

Nao Suzuki,
St. Marianna University
School of Medicine, Japan

Reviewed by:

Nalini Mahajan,
Mother & Child Hospital, India
Seido Takae,
St. Marianna University
School of Medicine, Japan

*Correspondence:

Claus Yding Andersen
claus.yding.andersen@regionh.dk

Specialty section:

This article was submitted
to Reproduction,
a section of the journal
Frontiers in Endocrinology

Received: 28 March 2017

Accepted: 17 May 2017

Published: 02 June 2017

Citation:

Borgbo T, Klučková H, Macek M Sr.,
Chrudimska J, Kristensen SG,
Hansen LL and Andersen CY (2017)
The Common Follicle-Stimulating
Hormone Receptor (FSHR) Promoter
Polymorphism FSHR –29G > A
Affects Androgen Production in
Normal Human Small Antral Follicles.
Front. Endocrinol. 8:122.
doi: 10.3389/fendo.2017.00122

Follicle-stimulating hormone receptors (FSHRs) are almost exclusively expressed on granulosa cells, and FSH action is probably most clearly reflected in intrafollicular hormone milieu of antral follicles. Little is known about the possible effects of the common single nucleotide polymorphism (SNP) FSHR –29G > A (rs1394205) on hormonal conditions in human small antral follicles (hSAFs) obtained from women in the natural menstrual cycle. This study investigated the follicle fluid (FF) concentrations of anti-Müllerian hormone, estradiol, progesterone, androstenedione, and testosterone in hSAF in relation to the different genotypes of FSHR –29G > A. FF from 362 follicles was collected in 95 women undergoing fertility preservation, who did not suffer from a disease that directly affected ovarian function. The testosterone levels of the minor A/A genotype were significantly increased compared to the A/G and the G/G genotype. Furthermore, significantly reduced androstenedione levels were observed for the G/G genotype, as compared to the A/G genotype, while the other hormones did not show statistical significant differences. In conclusion, the androgen levels of hSAF were significantly elevated in the minor SNP genotype in the FSHR promoter polymorphism FSHR –29G > A.

Keywords: follicle fluid, follicle-stimulating hormone receptor –29G > A, follicle-stimulating hormone receptor polymorphisms, human small antral follicles, rs1394205

INTRODUCTION

FSH is instrumental in regulating ovarian activity and exerts a myriad of effects on follicles in all developmental stages through follicle-stimulating hormone receptors (FSHRs), which are almost exclusively expressed on granulosa cells (GCs). The number of FSHR proteins expressed on each GC change profoundly throughout follicular development. Low but measurable FSHR expression has been observed in human primordial follicles (1, 2), while maximal expression occurs in the antral stage declining toward ovulation (3). FSH signal transduction is effected *via* different pathways of which increase in intracellular cAMP production leading to activation of protein kinase A and transcription of a several downstream genes is one of the major ones (4–7). It is, however, currently unknown how FSHR density impacts on function and signal transduction.

Follicle-stimulating hormone receptors exist as a number of different genetic variants, i.e., polymorphisms, which may affect signal transduction. Three common single nucleotide polymorphisms (SNPs) have been identified in the FSHR gene: rs6165 (c.919A > G, p.Thr307Ala, FSHR 307), rs6166 (c.2039A > G, p.Asn680Ser, FSHR 680), and rs1394205 (c. -29G > A, FSHR -29G > A) (8–14).

The two most studied SNPs, FSHR 307, and FSHR 680, both reside on exon 10 in linkage disequilibrium. Studies have focused on the intrafollicular hormone milieu in human small antral follicles (hSAFs) (15) and especially their potential impact on ovarian stimulation (OS) have received a considerable interest (8–14, 16, 17). Although single studies has found an association to basic levels of FSH, the length of menstrual cycles, as well as the consumption of FSH used in OS and treatment outcome, no clear cut effects have been found (11, 14, 17–23). However, recently, Lindgreen and co-workers found a significant association to pregnancy rates especially in combination with a LH SNP (24).

The third common SNP FSHR -29G > A is located in the promoter region of FSHR and has been studied to a lesser extent. So far, studies have associated FSHR -29G > A to changes in FSHR gene expression levels, in which homozygote carriers of the minor allele (i.e., the A/A genotype) showed a reduced FSHR gene expression, compared to homozygote carriers of the major allele (i.e., the G/G genotype) (9, 10, 19). Furthermore, Desai and co-workers observed that carriers of the minor genotype required significantly higher amounts of exogenous FSH during OS, when compared to the G/G and G/A genotypes (9).

Potential differences in the signal transduction of the different FSHR polymorphisms is probably best studied in FF, as the concentrations of locally produced substances such as sex steroids and anti-Müllerian hormone (AMH) often shown concentrations up to a thousand times higher than in circulation (25–27). The aim of this study was to evaluate the impacts of the FSHR -29G > A genotype on the hormone profile in FF from hSAF.

MATERIALS AND METHODS

Sample Material

In total, 362 follicles were collected from 95 women, with a mean age of 28.5 years (ranging from 15 to 38 years). The number of follicles analyzed from each woman ranged from 1 to 10 (mean 3.8 follicles/patient). Follicle diameters ranged from 3 to 13 mm, as calculated from the aspirated volume, assuming a spherical structure of the follicle. All follicles were obtained from surplus ovarian tissue from women undergoing fertility preservation, having ovarian tissue cryopreserved at Laboratory of Reproductive Biology, Rigshospitalet, Denmark. The diagnosis of women was in all cases unrelated to the ovary including polycystic ovary syndrome. The fertility preservation procedure involved excision of one entire ovary from which individual visible antral follicles were aspirated with a 23G needle attached to a syringe. The collection of FF had no effect on the fertility preservation procedure. In all cases, the ovary had overall gross normal appearance.

As the intrafollicular hormone milieu in human small antral follicles (hSAFs) are independent of circulating levels of gonadotropins

and have no major impact on the serum hormone levels (15), serum hormone levels were not included in this study. The follicles were collected at various times during the menstrual cycle, as the dynamics of the hSAF are similar throughout the menstrual cycle (15, 28).

Parts of this material has been used in previous studies (3, 15, 27, 29, 30).

Ethical Approval

The study was approved by the ethical committee of the municipalities of Copenhagen and Frederiksberg (H-2-2011-044). Oral information and written informed consent was obtained from research participants and from parents of minors.

DNA Extraction

Seventy-two patients were genotyped using DNA extracted from approximately 25 mg ovarian tissue, using the DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany), following the manufacturers protocol. The DNA samples were subsequently stored at -20°C. The remaining 23 patients were genotyped from DNA extracted from aspirated GC, as no ovarian tissue was available for these patients. The DNA was extracted from the GC samples using Tri Reagent (Sigma-Aldrich, St. Louis, MO, USA), with slight modifications to the protocol as previously described (30).

Genotyping of the FSHR -29G > A Polymorphism

The samples were genotyped using CADMA-based genotyping as previously published with slight modifications to the PCR master-mix (30). In brief, the final reaction mix consisted of 2 µl template DNA (10 ng/µl), 5 µl LightCycler® 480 High Resolution Melting Master (Roche Diagnostics, Ropkreuz, Switzerland), 1.2 µl MgCl₂ (25 µM, Roche Diagnostics, Ropkreuz, Switzerland), 0.2 µl wild-type primer 5'-TATGCATCCGTCACCTGAGTTCTTC-3' (10 µM), 0.2 µl mutation primer 5'-TATGCATCTATCCACA TGATTCTTT-3' (10 µM), 0.2 µl common reverse primer 5'-GAGGTTTTTCTCTGCAAATGCAG-3' (10 µM), and finally adding 1.2 µl ddH₂O to a final volume of 10 µl.

Hormone Measurements

The concentration of AMH, estradiol, progesterone, androstenedione, and testosterone were measured using commercially available ELISA assays. AMH was measured using the Ultra Sensitive AMH/MIS Elisa kit (AI-105-i, AnshLabs, Webster, TX, USA), according to the manufacturer's protocol, with a 1:300 dilution of the FF samples using the supplied assay buffer. Estradiol, progesterone, androstenedione, and testosterone were initially measured using commercially available RIA kits (DSL-43100, DSL-3400, DSL-3800, DSL-4000; DSL, Webster, TX, USA). However, during the course of sample collection, the RIA assays became unavailable and it was necessary to switch to ELISA assays (NovaTec Immundiagnostica, Dietzenbach, Germany, DNOV 002, 003, 006 and 008, respectively).

Due to limited sample material, it was not possible to re-analyze the FF analyzed with RIA. The RIA kit was used to analyze between 11 and 15% of the FF samples in this study; progesterone

($n = 39/336$, 12%), estradiol ($n = 37/337$, 11%), testosterone ($n = 47/338$, 14%), and androstenedione ($n = 50/345$, 15%). Based on highly significant linear correlations between the results from the two types of assays (correlation coefficients: estradiol $r = 0.99$, progesterone $r = 0.93$, testosterone $r = 0.91$, and androstenedione $r = 0.91$), mathematical equations was used for transforming data from RIA into ELISA, as previously published (3). The remaining follicles were analyzed using NovaTech ELISA assays, according to the manufacturer's protocol, using in-house prepared steroid-free serum for FF dilution.

Statistical Analysis

The statistical analysis was performed using STATA version 13.1 (STATACorp LP, USA). The hormones were log- or square root-transformed in order to approximate normal distributions. A one-level mixed effects model was used for the statistical analysis, including follicle size as covariant and patients as random effect, in order to take into account that GC function varies with follicle size, and that follicles from one patient may be more similar than follicles from different patients. P -values ≤ 0.05 were considered statistically significant.

In this study, we have included follicles ranging from 3 to 13 mm in diameter. The intrafollicular hormone production is dependent on the developmental stage and size of the follicles, and a shift in hormone profiles is expected when follicles reach 810 mm when selection for dominance is initiated. In the follicle cohort of this study; however, no differences were found in the statistics, when excluding follicles above 8 mm (data not published).

RESULTS

The distribution between the three FSHR $-29G > A$ genotypes (i.e., A/A, A/G, and G/G allele) was 14, 37, and 49%, respectively. This frequency distribution compares well to what have been reported from other countries (9, 13, 31).

The FF hormone concentrations are summarized in **Table 1** and **Figure 1**. Overall, the FSHR $-29G > A$ genotype was found to be significantly associated to the androgen levels in hSAF. Significantly increased testosterone levels were observed for the A/A genotype as compared to both the A/G ($P = 0.02$) and G/G ($P = 0.02$) genotype. Likewise, significantly lower androstenedione levels were observed for the G/G genotype as compared to the A/G genotype ($P = 0.03$), however, with no significant differences observed between the A/A and the A/G genotype.

No statistically significant differences were observed for AMH, estradiol, or progesterone.

DISCUSSION

On a relatively large sample size of more than 300 normal hSAF, the present study demonstrated that the minor genotype of the common SNP at position -29 of the FSHR promoter region, FSHR $-29G > A$, is significantly associated to augmented FF concentrations of androstenedione and testosterone. Androgens exert an important role in follicular development and in female reproduction, and the present results enforce that SNP's in the gonadotropin receptors affect the fine-tuned pituitary gonadal interactions. The present study was performed entirely on hSAF from women who did not receive any kind of exogenous stimulation in their natural menstrual cycle and it remains to be seen whether the supraphysiological concentrations of FSH as used is connection with OS override these subtle differences and wipe out any impact on the reproductive output of fertility treatment.

Androgen production takes place exclusively in the theca cells, which do not express FSHR and there is no direct effect of FSH on theca cells. However, in rats, it has been shown that FSH-induced paracrine factors from GCs enhance theca cell androgen synthesis (32). Treatment with FSH was found to significantly increase the thecal CYP17A1 mRNA levels, as well as the androgen production in response to LH (32). Smyth and co-workers found that inhibin enhanced the LH-induced androgen

TABLE 1 | Intrafollicular hormone levels grouped according to follicle-stimulating hormone receptor $-29G > A$ genotype mean \pm SEM.

Genotype	A/A			A/G			G/G		
	<i>n</i> follicle fluid (FF)	<i>n</i> (women)	Mean \pm SEM	<i>n</i> (FF)	<i>n</i> (women)	Mean \pm SEM	<i>n</i> (FF)	<i>n</i> (women)	Mean \pm SEM
Anti-Müllerian hormone (pmol/l)	48	9	8,814 \pm 746	133	37	6,266 \pm 441	166	46	7,832 \pm 510
Progesterone (mol/l)	49	10	165 \pm 48	123	37	168 \pm 26	164	46	160 \pm 19
Estradiol (E_2) (nmol/l)	48	10	130 \pm 33	123	37	217 \pm 31	166	47	142 \pm 22
Testosterone (T) (nmol/l)	49	10	386 \pm 32 ^{1,2}	125	37	230 \pm 14 ^{*1}	164	45	236 \pm 13 ^{*2}
Androstenedione (nmol/l)	49	10	2,157 \pm 188	125	37	1,919 \pm 104 ^{*3}	171	47	1,521 \pm 87 ^{*3}
Age \pm SD (years)		10	29.6 \pm 5.8		37	28.8 \pm 5.7		45	28.0 \pm 5.9
Follicle diameter (mm)	50	10	5.3 \pm 0.2	135	37	6.1 \pm 0.1	177	48	5.6 \pm 0.1

*Statistical significances.

Testosterone: ¹ $P = 0.02$, ² $P = 0.02$.

Androstenedione: ³ $P = 0.03$.




FIGURE 1 | Follicular fluid hormone profiles grouped according to follicle stimulating hormone receptor $-29G > A$ genotype. The figure displays the Mean \pm SEM of the intrafollicular hormone levels. Statistically significant differences in hormone levels are observed for androstenedione, and testosterone. In addition, discrete genotype-dependent differences are observed for Anti-Müllerian hormone and estradiol in an inverse manner, although not statistically significant.

production and suggested that FSH-induced paracrine factors produced by the GCs (e.g., inhibin-B or IGF2) could augment theca cell androgen output. More recently, we found a statistical significant association between inhibin-B and the androgen concentration in hSAF (33, 34). In the present study, inhibin-B was only measured in a small fraction of FF that did not allow a valid analysis.

Although the minor allele of FSHR $-29G > A$ has been associated with reduced FSHR gene expression levels (9, 10, 19) and, therefore, would be expected to result in less paracrine androgen stimulation, it is not known how and if this specific SNP affect inhibin-B secretion. Especially, the inhibin α -gene has been shown to be upregulated by acidic FSH isoforms in contrast to almost any other FSH-inducible GC substance, where the less acidic isoforms are more potent (35). It could, therefore, also be speculated that the minor allele of FSHR $-29G > A$ had a similar effect. It will, therefore, be of interest to evaluate the levels of inhibin-B and inhibin-A in FF from hSAF in order to evaluate whether levels of inhibin is also associated to this SNP.

The paracrine effects of androgens-enhancing GC FSHR expression has been documented in several studies (36, 37). Thus, GC from follicles obtained from women having the minor allele A/A genotype, which showed augmented androgen FF concentration, would be expected to show enhanced FSHR expression. The increased FSHR expression is expected to translate into an augmented aromatase expression. However,

previous studies showed that the minor allele group had a reduced FSHR expression (9, 10, 19), which may explain why levels of estradiol are not significantly enhanced. It would be relevant to evaluate both the FSHR and LHR expression of the corresponding GCs to substantiate the effects of androgens in a future study.

In conclusion, this study shows that the minor A/A genotype at position -29 of FSHR is associated with an altered FF milieu, resulting in significant increased androgen levels.

ETHICS STATEMENT

This study was carried out in accordance with the recommendations of the ethical committee of the municipalities of Copenhagen and Frederiksberg with written informed consent from all subjects. All subjects gave written informed consent in accordance with the Declaration of Helsinki. The protocol was approved by the ethical committee of the municipalities of Copenhagen and Frederiksberg (journal number; H-2-2011-044).

AUTHOR CONTRIBUTIONS

TB, CA, and MM designed the study and wrote the manuscript. TB performed the primer design for the CADMA-based FSHR assays, and performed the CADMA-based FSHR $-29G > A$ genotyping. LH provided validating sequencing results of the CADMA-based FSHR assays. HK and JC performed validating

FSHR $-29G > A$ HRM genotyping. SK and CA collected follicular fluids, granulosa cells, and ovarian tissue for the study. TB performed the analysis and interpretation of the data, and prepared figures and tables for the manuscript. LH helped finalizing the manuscript.

ACKNOWLEDGMENTS

The authors would like to acknowledge Marjo Westerdahl and Jane Alrø Bøtkjaer for performing parts of the ELISA analysis for this study. The authors would like to acknowledge Bo Martin Bibby for consultancy regarding the statistical analysis of the data. TB

REFERENCES

- Kristensen SG, Ebbesen P, Andersen CY. Transcriptional profiling of five isolated size matched stages of human preantral follicles. *Mol Cell Endocrinol* (2015) 401:189–201. doi:10.1016/j.mce.2014.12.012
- Oktay K, Briggs D, Gosden RG. Ontogeny of follicle-stimulating hormone receptor gene expression in isolated human ovarian follicles. *J Clin Endocrinol Metab* (1997) 82:3748–51. doi:10.1210/jcem.82.11.4346
- Jeppesen JV, Kristensen SG, Nielsen ME, Humaidan P, Dal Canto M, Fadini R, et al. LH-receptor gene expression in human granulosa and cumulus cells from antral and preovulatory follicles. *J Clin Endocrinol Metab* (2012) 97:E1524–31. doi:10.1210/jc.2012-1427
- Escamilla-Hernandez R, Little-Ihrig L, Orwig KE, Yue J, Chandran U, Zeleznik AJ. Constitutively active protein kinase A qualitatively mimics the 294 effects of follicle-stimulating hormone on granulosa cell differentiation. *Mol Endocrinol* (2008) 22:1842–52. doi:10.1210/me.2008-0103
- Lavoie HA, King SR. Transcriptional regulation of steroidogenic genes: STARD1, CYP11A1 and HSD3B. *Exp Biol Med* (Maywood) (2009) 234:880–907. doi:10.3181/0903-MR-97
- Menon KJM, Menon B. Structure, function and regulation of gonadotropin receptors – a perspective. *Mol Cell Endocrinol* (2012) 356:88–97. doi:10.1016/j.mce.2012.01.021
- Priyanka S, Medhamurthy R. Characterization of cAMP/PKA/CREB signaling cascade in the bonnet monkey corpus luteum: expressions of inhibin-alpha and StAR during different functional status. *Mol Hum Reprod* (2007) 13:381–90. doi:10.1093/molehr/gam015
- Casarini L, Pignatti E, Simoni M. Effects of polymorphisms in gonadotropin and gonadotropin receptor genes on reproductive function. *Rev Endocr Metab Disord* (2011) 12:303–21. doi:10.1007/s11154-011-9192-2
- Desai SS, Achrekar SK, Pathak BR, Desai SK, Mangoli VS, Mangoli RV, et al. Follicle-stimulating hormone receptor polymorphism (G-29A) is associated with altered level of receptor expression in granulosa cells. *J Clin Endocrinol Metab* (2011) 96:2805–12. doi:10.1210/jc.2011-1064
- Nakayama T, Kuroi N, Sano M, Tabara Y, Katsuya T, Ogihara T, et al. Mutation of the follicle stimulating hormone receptor gene 5'-untranslated region associated with female hypertension. *Hypertension* (2006) 48:512–8. doi:10.1161/01.HYP.0000233877.84343.d7
- Perez Mayorga M, Gromoll J, Behre HM, Gassner C, Nieschlag E, Simoni M. Ovarian response to follicle-stimulating hormone (FSH) stimulation depends on the FSH receptor genotype. *J Clin Endocrinol Metab* (2000) 85:3365–9. doi:10.1210/jcem.85.9.6789
- Themmen APN, Huhtaniemi IT. Mutations of gonadotropins and gonadotropin receptors: elucidating the physiology and pathophysiology of pituitary-gonadal function. *Endocr Rev* (2000) 21:551–83. doi:10.1210/edrv.21.5.0409
- Wunsch A, Ahda Y, Banaz-Yaşar F, Sonntag B, Nieschlag E, Simoni M, et al. Single-nucleotide polymorphisms in the promoter region influence the expression of the human follicle-stimulating hormone receptor. *Fertil Steril* (2005) 84:446–53. doi:10.1016/j.fertnstert.2005.02.031
- Yan Y, Gong Z, Zhang L, Li Y, Li X, Zhu L, et al. Association of follicle-stimulating hormone receptor polymorphisms with ovarian response in Chinese women: a prospective clinical study. *PLoS One* (2013) 8:e78138. doi:10.1371/journal.pone.0078138
- Borgbo T, Jeppesen JV, Lindgren I, Lundberg Giwerzman Y, Hansen LL, Yding Andersen C. Effect of the FSH receptor single nucleotide polymorphisms (FSHR 307/680) on the follicular fluid hormone profile and the granulosa cell gene expression in human small antral follicles. *Mol Hum Reprod* (2015) 21:255–61. doi:10.1093/molehr/gau106
- Casarini L, Moriondo V, Marino M, Adversi F, Capodanno F, Grisolia C, et al. FSHR polymorphism p.N680S mediates different responses to FSH in vitro. *Mol Cell Endocrinol* (2014) 393:83–91. doi:10.1016/j.mce.2014.06.013
- Greb RR, Behre HM, Simoni M. Pharmacogenetics in ovarian stimulation – current concepts and future options. *Reprod Biomed Online* (2005) 11:589–600. doi:10.1016/S1472-6483(10)61167-4
- Boudjenah R, Molina-Gomes D, Torre A, Bergere M, Bailly M, Boitrelle F, et al. Genetic polymorphisms influence the ovarian response to rFSH stimulation in patients undergoing in vitro fertilization programs with ICSI. *PLoS One* (2012) 7:e38700. doi:10.1371/journal.pone.0038700
- Jun JK, Yoon JS, Ku S-Y, Choi YM, Hwang KR, Park SY, et al. Follicle-stimulating hormone receptor gene polymorphism and ovarian responses to controlled ovarian hyperstimulation for IVF-ET. *J Hum Genet* (2006) 51:665–70. doi:10.1007/s10038-006-0005-5
- Altmäe S, Hovatta O, Stavreus-Evers A, Salumets A. Genetic predictors of controlled ovarian hyperstimulation: where do we stand today? *Hum Reprod Update* (2011) 17:813–28. doi:10.1093/humupd/dmr034
- de Castro F, Ruiz R, Montoro L, Pérez-Hernández D, Sánchez-Casas Padilla E, Real LM, et al. Role of follicle-stimulating hormone receptor Ser680Asn polymorphism in the efficacy of follicle-stimulating hormone. *Fertil Steril* (2003) 80:571–6. doi:10.1016/S0015-0282(03)00795-7
- Klinkert ER, te Velde ER, Weima S, van Zandvoort PM, Hanssen RGJM, Nilsson PR, et al. FSH receptor genotype is associated with pregnancy but not with ovarian response in IVF. *Reprod Biomed Online* (2006) 13:687–95. doi:10.1016/S1472-6483(10)60660-8
- Loutradis D, Theofanakis C, Anagnostou E, Mavrogianni D, Partsinevelos GA. Genetic profile of SNP(s) and ovulation induction. *Curr Pharm Biotechnol* (2012) 13:417–25. doi:10.2174/138920112799361954
- Lindgren I, Bååth M, Uvebrant K, Dejmeck A, Kjaer L, Henic E, et al. Combined assessment of polymorphisms in the LHCGR and FSHR genes predict chance of pregnancy after in vitro fertilization. *Hum Reprod* (2016) 31(3):672–83. doi:10.1093/humrep/dev342
- Andersen CY. Characteristics of human follicular fluid associated with successful conception after in vitro fertilization. *J Clin Endocrinol Metab* (1993) 77:1227–34. doi:10.1210/jcem.77.5.7521343
- Andersen CY, Schmidt KT, Kristensen SG, Rosendahl M, Byskov AG, Ernst E. Concentrations of AMH and inhibin-B in relation to follicular diameter in normal human small antral follicles. *Hum Reprod* (2010) 25:1282–7. doi:10.1093/humrep/deq019
- Jeppesen JV, Anderson RA, Kelsey TW, Christiansen SL, Kristensen SG, Jayaprakasan K, et al. Which follicles make the most anti-Müllerian hormone in humans? Evidence for an abrupt decline in AMH production at the time of follicle selection. *Mol Hum Reprod* (2013) 19:519–27. doi:10.1093/molehr/gat024
- Kuang Y, Hong Q, Chen Q, Lyu Q, Ai A, Fu Y, et al. Luteal-phase ovarian stimulation is feasible for producing competent oocytes in women undergoing in vitro fertilization/intracytoplasmic sperm injection treatment, with optimal

was granted a Ph.D. scholarship from Health, Aarhus University, Denmark.

FUNDING

The study was financial supported from University of Aarhus, the Novo Nordic Foundation, Sophus Carl Emil Friis and wife Olga Doris Friis' foundation, the Lundbeck Foundation, and the University Hospital of Copenhagen. MM was supported by the Czech Ministry of Health Conceptual Development of Research Organization (University Hospital Motol, Prague, Czech Republic; grants 00064203, NT/13770-4, and CZ.2.16/3.1.00/24022OPPK).

- pregnancy outcomes in frozen-thawed embryo transfer cycles. *Fertil Steril* (2014) 101:105–11. doi:10.1016/j.fertnstert.2013.09.007
29. Borgbo T, Macek M, Chrudimska J, Jeppesen JV, Hansen LL, Andersen CY. Size matters: associations between the androgen receptor CAG repeat length and the intrafollicular hormone milieu. *Mol Cell Endocrinol* (2016) 419:12–7. doi:10.1016/j.mce.2015.09.015
 30. Borgbo T, Sommer Kristensen L, Lindgren I, Yding Andersen C, Hansen LL, Kristensen LS. Genotyping common FSHR polymorphisms based on competitive amplification of differentially melting amplicons (CADMA). *J Assist Reprod Genet* (2014) 31:1427–36. doi:10.1007/s10815-014-0329-6
 31. Simoni M, Casarini L. Mechanisms in endocrinology: genetics of FSH action: a 2014-and beyond view. *Eur J Endocrinol* (2014) 170:R91–107. doi:10.1530/EJE-13-0624
 32. Smyth CD, Miró F, Whitelaw PF, Howles CM, Hillier SG. Ovarian thecal/interstitial androgen synthesis is enhanced by a follicle-stimulating hormone-stimulated paracrine mechanism. *Endocrinology* (1993) 133:1532–8. doi:10.1210/endo.133.4.8404591
 33. Yding Andersen C, Rosendahl M, Byskov AG. Concentration of anti-Müllerian hormone and inhibin-B in relation to steroids and age in follicular fluid from small antral human follicles. *J Clin Endocrinol Metab* (2008) 93:2344–9. doi:10.1210/jc.2007-2342
 34. Yding Andersen C. Inhibin-B secretion and FSH isoform distribution may play an integral part of follicular selection in the natural menstrual cycle. *Mol Hum Reprod* (2017) 23(1):16–24. doi:10.1093/molehr/gaw070
 35. Timossi CM, Barrios-de-Tomasi J, González-Suárez R, Arranz MC, Padmanabhan V, Conn PM, et al. Differential effects of the charge variants of human follicle stimulating hormone. *J Endocrinol* (2000) 165:193–205. doi:10.1677/joe.0.1650193
 36. Weil S, Vendola K, Zhou J, Bondy CA. Androgen and follicle-stimulating hormone interactions in primate ovarian follicle development. *J Clin Endocrinol Metab* (1999) 84:2951–6. doi:10.1210/jcem.84.8.5929
 37. Nielsen ME, Rasmussen IA, Kristensen SG, Christensen ST, Møllgård K, Wreford Andersen E, et al. In human granulosa cells from small antral follicles, androgen receptor mRNA and androgen levels in follicular fluid correlate with FSH receptor mRNA. *Mol Hum Reprod* (2011) 17:63–70. doi:10.1093/molehr/gaq073

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer ST and handling Editor declared their shared affiliation, and the handling Editor states that the process nevertheless met the standards of a fair and objective review.

Copyright © 2017 Borgbo, Klučková, Macek, Chrudimska, Kristensen, Hansen and Andersen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.